arXiv:astro-ph/0404208v1 9 Apr 2004

BL LAC EVOLUTION REVISITED

ANNA WOLTER, FRANCESCA CAVALLOTTI
INAF- Osservatorio Astronomico di Brera
Via Brera, 28 20121 Milano, ITALY
E-mail: anna@brera.mi.astro.it; cavallot@brera.mi.astro.it

JOHN T. STOCKE
Center for Astrophysics and Space Astronomy
University of Colorado, Boulder, CO 80309-0389, USA

TRAVIS RECTOR
University of Alaska
3211 Providence Dr., BMB 212, Anchorage, AK 99508, USA

BL Lac objects are an elusive and rare class of active galactic nuclei. For years their evolutionary behavior has appeared inconsistent with the trend observed in the population of AGN at large. The so-called "negative" evolution implies that BL Lacs were either less or fainter in the past. This effect is stronger for BL Lacs selected in X-ray surveys. We have investigated if one of the selection criteria, namely the flat-radio spectrum (imposed on the Radio-selected but not on the X-ray-selected samples), might explain the different evolutionary trend.

1. BL Lacs samples and evolution

Radio selected and X-ray selected samples of BL Lac objects are created through various selections and show different evolutionary properties as shown in Table 1. It is evident from the Table indications that only samples that impose the α_r (flat radio spectrum) criterion show positive evolution (vs. negative or no-evolution). We want to analyze whether this selection criterion is a primary cause for the different evolution measured in different sample. We obtained simultaneous data at the Very Large Array (VLA[a]) at 20, 6 and 3.6 cm (C conf.) for a (radio) flux limited sample of 26 sources in the EMSS ([1],[2]) with the purpose of studying the distribution of α_r in a complete unbiased sample. A detailed analysis is available in [3].

---

[a] The VLA is a facility of the National Radio Astronomy Observatory. The National Radio Astronomy Observatory (NRAO) is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

Table 1. Summary of BL Lacs samples (their size), measure of evolution and selection criteria applied

Samples (#)	Evolution is:	Selection criteria applied
X-ray selected		
EMSS (40 objects)	negative	f_X
Radio selected		
1Jy (34 objects)	*positive*/null	f_r - mag - α_r
Radio & X-ray selected		
RGB (33 objects)	negative	f_X - f_r - mag
DXRBS (30 objects)	*positive*	f_X - f_r - α_r
REX (55 in XB-REX)	negative/null	f_X - f_r - mag
HRX (77 objects)	negative	f_X - f_r
Others		
Sedentary (58 objs)	negative	f_X - f_r - mag -α_{ro} α_{rx} - (HR)
[PG - (6 objs)]	unknown	UV excess

Note: EMSS: Rector et al. 2000 AJ 120 1626; 1Jy: Stickel et al. 1991 ApJ 374 431; RGB: Laurent-Muehleisen et al. 1993 AJ 106 875; DXRBS: Landt et al. 2001 MNRAS 323 757; REX: Caccianiga et al. 2002 ApJ 566 181; HRX: Beckmann et al. 2003 A&A 411 327; Sedentary: Giommi et al. 1999 MNRAS 310 65; PG: Schmidt & Green 1983, ApJ 269 352.

We find that a large fraction of BL Lacs show a steep spectrum (15-40% depending on conditions of observations). These objects are missed in radio surveys that need an α_r selection criterion to avoid the bulk of radio galaxies. However, on average, steep spectrum sources have similar properties to flat spectrum sources. We conclude that the spectral selection is not responsible for the different evolution properties of RBL and XBL.

Acknowledgments

This work has received partial financial support from the Italian Space Agency (ASI) and MIUR.

References

1. Gioia I.M., et al. 1990, ApJS, 72, 567.
2. Stocke et al. 1991, ApJS, 76, 813.
3. Cavallotti et al. 2004, to be published in A&A.